

# Exercise Sheet 13

Discussed on 21.07.2021

**Definition.** Let  $K$  be a field,  $V$  a  $K$ -vector space and  $P: V \rightarrow K$  a map. We say that  $P$  is *polynomial* if for any  $n \geq 0$  and any vectors  $v_1, \dots, v_n \in V$  the map

$$K^n \rightarrow K, \quad (x_1, \dots, x_n) \mapsto P(x_1v_1 + \dots + x_nv_n)$$

is given by a polynomial, i.e. by an element of  $K[T_1, \dots, T_n]$ .

**Lemma.** Let  $K$  be an infinite field,  $V$  a  $K$ -vector space and  $P: V \rightarrow K$  a map. Then  $P$  is *polynomial* if and only if for all  $v, w \in V$  the map

$$K \rightarrow K, \quad x \mapsto P(v + xw)$$

is given by a polynomial.

**Problem 1.** Let  $X$  be an abelian variety of dimension  $g$  over some field  $k$ .

- (a) Let  $\varphi, \psi \in \text{End}(X)$  and let  $L$  be a line bundle on  $X$ . Show that there are line bundles  $L_0, L_1, L_2$  on  $X$  such that for all  $n \in \mathbb{Z}$  we have

$$(\psi + n\varphi)^*L = L_0 \otimes L_1^n \otimes L_2^{n(n-1)/2}.$$

- (b) Show that  $\text{deg}: \text{End}(X) \rightarrow \mathbb{Z}$  extends to a polynomial function on  $\text{End}^0(X) = \text{End}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$  (here  $\text{deg } \varphi = 0$  if  $\varphi$  is not surjective).
- (c) For every prime  $\ell$ , define the  $\ell$ -adic Tate module  $T_\ell X$  of  $X$ . Show that the natural map

$$\mathbb{Z}_\ell \otimes_{\mathbb{Z}} \text{End}(X) \hookrightarrow \text{End}(T_\ell X)$$

is injective. Deduce that  $\text{End}(X)$  has  $\text{rank} \leq 4g^2$ .

*Hint:* Apply the same proof strategy as for elliptic curves.